

New York State Department of Health Adult Immunization Schedule

Vaccine	19-49 Years	50-64 Years	65 Years or Older
Influenza	You need one dose yearly if: you are a health care worker; you have a chronic health problem;* you come into close contact with individuals at risk for complications of influenza;* or you just want to avoid getting influenza.	You need one dose yearly.	You need one dose yearly.
Pneumococcal	You need one to two doses if you have certain chronic medical conditions.*	You need one to two doses if you have certain chronic medical conditions and you have not previously received this vaccine.*	You need one dose at 65 years or older.
Tetanus, Diphtheria, Pertussis (Td, Tdap)	<p>If you've never had at least three shots containing tetanus and diphtheria, you need to get them now.</p> <p>In addition, all adults need another Td (tetanus and diphtheria) shot, or booster, every 10 years.</p> <p>If you're younger than 65, and as an adult you haven't had a vaccine that contains pertussis (whooping cough), one of the doses that you receive should have pertussis vaccine in it. This vaccine is known as Tdap.</p> <p>Be sure to consult your health care professional if you have a deep or dirty wound. You may need a dose at that time.</p>	<p>If you've never had at least three shots containing tetanus and diphtheria, you need to get them now.</p> <p>In addition, all adults need another Td (tetanus and diphtheria) shot, or booster, every 10 years.</p> <p>If you're younger than 65, and as an adult you haven't had a vaccine that contains pertussis (whooping cough), one of the doses that you receive should have pertussis vaccine in it. This vaccine is known as Tdap.</p> <p>Be sure to consult your health care professional if you have a deep or dirty wound. You may need a dose at that time.</p>	<p>If you've never had at least three shots containing tetanus and diphtheria, you need to get them now.</p> <p>In addition, all adults need another Td (tetanus and diphtheria) shot, or booster, every 10 years.</p> <p>Be sure to consult your health care professional if you have a deep or dirty wound. You may need a dose at that time.</p>
Hepatitis B (HepB)	<p>You need this vaccine if you have a specific risk factor for hepatitis B virus infection* or if you simply wish to be protected from this disease.</p> <p>This vaccine is given as a three-dose series.</p>	<p>You need this vaccine if you have a specific risk factor for hepatitis B virus infection* or you simply wish to be protected from this disease.</p> <p>This vaccine is given as a three-dose series.</p>	<p>You need this vaccine if you have a specific risk factor for hepatitis B virus infection* or you simply wish to be protected from this disease.</p> <p>This vaccine is given as a three-dose series.</p>

***Consult your healthcare professional to determine your level of risk for infection and your need for this vaccine.**

Source: Immunization Action Coalition www.immunize.org

New York State Department of Health Adult Immunization Schedule

Vaccine	19-49 Years	50-64 Years	65 Years or Older
Hepatitis A (HepA)	You need this vaccine if you have a specific risk factor for hepatitis A virus infection* or you simply wish to be protected from this disease. This vaccine is usually given as two doses, 6–18 months apart.	You need this vaccine if you have a specific risk factor for hepatitis A virus infection* or you simply wish to be protected from this disease. This vaccine is usually given as two doses, 6–18 months apart.	You need this vaccine if you have a specific risk factor for hepatitis A virus infection* or you simply wish to be protected from this disease. This vaccine is usually given as two doses, 6-18 months apart.
Human Papillomavirus (HPV)	You need this vaccine if you are a woman who is 26 years or younger. This vaccine is given as a three-dose series.		
Measles, Mumps, Rubella (MMR)	You need at least one dose of MMR if you were born in 1957 or later. You also may need a second dose.*	You need at least one dose of MMR if you were born in 1957 or later. You also may also need a second dose.*	
Varicella (Chickenpox)	If you've never had chickenpox, or you were never vaccinated, you should complete a two-dose series. If you've been vaccinated but you received only one dose, you should get a second dose.	If you've never had chickenpox, or you were never vaccinated, you should complete a two-dose series. If you've been vaccinated but you received only one dose, you should get a second dose.	If you've never had chickenpox, or you were never vaccinated, you should complete a two-dose series. If you've been vaccinated but you received only one dose, you should get a second dose.
Meningococcal	If you are going to college and you plan to live in a dormitory, you should get vaccinated against meningococcal disease. People with certain medical conditions also should receive this vaccine.*	People with certain medical conditions should receive this vaccine.*	People with certain medical conditions should receive this vaccine.*
Zoster (Shingles)		If you are 60 years or older, you should get this vaccine.	If you are 60 years or older, you should get this vaccine.
Do you travel outside the United States?	If so, you may need additional vaccines. The Centers for Disease Control and Prevention (CDC) operates an international traveler's health information line. Call 1-877-394-8747 or visit CDC's Web site at www.cdc.gov/travel for information about your destination. You may also consult a travel clinic or your health care professional. Source: Immunization Action Coalition www.immunize.org		
*Consult your healthcare professional to determine your level of risk for infection and your need for this vaccine.			